

Procedure Code Set General Equivalence Mappings

ICD-10-PCS to ICD-9-CM and ICD-9-CM to ICD-10-PCS

2009 Version

Documentation and User's Guide

Preface

Purpose and Audience

This document accompanies the 2009 update of the Centers for Medicare and Medicaid Studies (CMS) public domain code reference mappings of the ICD-10 Procedure Code System (ICD-10-PCS) and the International Classification of Diseases 9th Revision (ICD-9-CM) Volume 3. The purpose of this document is to give readers the information they need to understand the structure and relationships contained in the mappings so they can use the information correctly. The intended audience includes but is not limited to professionals working in health information, medical research and informatics. General interest readers may find section 1 useful. Those who may benefit from the material in both sections 1 and 2 include clinical and health information professionals who plan to directly use the mappings in their work. Software engineers and IT professionals interested in the details of the file format will find this information in Appendix A.

Document Overview

For readability, ICD-9-CM is abbreviated “I-9,” and ICD-10-PCS is abbreviated “PCS.” The network of relationships between the two code sets described herein is named the General Equivalence Mappings (GEMs).

- **Section 1** is a general interest discussion of mapping as it pertains to the GEMs. It includes a discussion of the difficulties inherent in translating between two coding systems of different design and structure. The specific conventions and terms employed in the GEMs are discussed in more detail.
- **Section 2** contains detailed information on how to use the GEM files for users who will be working directly with mapping applications—as coding experts, researchers, claims processing personnel, software developers, etc.
- The **Glossary** provides a reference list of the terms and conventions used—some unique to this document—with their accompanying definitions.
- **Appendix A** contains tables describing the technical details of the file formats, one for each of the two GEM files:
 - 1) ICD-9-CM to ICD-10-PCS (I-9 to PCS)
 - 2) ICD-10-PCS to ICD-9-CM (PCS to I-9)
- **Appendix B** contains a table listing the new I-9 procedure codes for October 1, 2008, and their corresponding entries in the I-9 to PCS GEM.

Section 1—Mapping and the GEMs

Translating Between the ICD-9 and ICD-10 Procedure Code Sets

Mappings between I-9 and I-10 attempt to find corresponding procedure codes between the two code sets, insofar as this is possible. Because the two systems are so different, translating between them the majority of the time can offer only a series of possible compromises rather than the mirror image of one code in the other code set.

A sentence translated from English to Chinese may not be able to capture the full meaning of the original because of fundamental differences in the structure of the language. Likewise, a mapping may not be able to seamlessly link the codes in one set to identical counterparts in the other code set, and this is especially true with I-9 and PCS. For these two procedure code sets, it is rare to find two corresponding descriptions that are identical in level of specificity and terminology used. This is understandable. Indeed, there would be little point in changing from the old system to the new system if the differences between the two, and the benefits available in the new system, were not significant.

There is no simple “crosswalk from I-9 to PCS” in the GEM files. A mapping that forces a simple correspondence—each I-9 code mapped only once—from the smaller, less detailed I-9 to the larger, more detailed PCS (a code set of entirely different design and scope) defeats the purpose of upgrading to PCS. It obscures the differences between the two code sets and eliminates any possibility of benefiting from the improvement in data quality that PCS offers. Instead of a simple crosswalk, the GEM files attempt to organize those differences in a meaningful way, by linking a code to all valid alternatives in the other code set from which choices can be made depending on the use to which the code is put.

It is important to understand the kinds of differences that need to be reconciled in linking coded data. The method used to reconcile those differences may vary, depending on whether the data is used for research, claims adjudication, or analyzing coding patterns between the two code sets; whether the desired outcome is to present an all-embracing look at the possibilities (one-to-many mapping) or to offer the one “best” compromise for the application (one-to-one mapping); whether the desired outcome is to translate existing coded data to their counterparts in the new code set (“forward mapping”) or to track newly coded data back to what they may have been in the previous code set (“backward mapping”), or any number of other factors. The scope of the differences varies, is complex, and cannot be overlooked if quality mapping and useful coded data are the desired outcomes. Several common types of differences between the code sets will be examined here in detail to give the reader a sense of the scope.

Procedure Codes and Differences in Structure

PCS is designed to avoid regional variants of code descriptions and “running out” of code capacity. It contains a standardized vocabulary of surgical concepts, body part terms, operative approaches, and so on, from which codes are built. For these reasons, translating between the two systems is often an “apples to oranges” enterprise.

The majority of the time, finding the “one correct” match in PCS coding concepts for a general I-9 concept is not possible. Because it is standardized, PCS contains code elements describing the precise objective of each coded procedure. Each of these concepts, called a “root operation” or “root type,” is defined in the system and can be used only when the procedure performed agrees with the root operation or type definition. Further, because a word used like “repair” as used in I-9 does not precisely identify a surgical method, the possible code alternatives in PCS must include all of the root operations that could have been performed. This means that any initial mapping between the systems must cast a wide net over possible equivalent options, options that can only be narrowed down when a specific clinical scenario, use case, or other application has been defined.

For example, an I-9 code description containing the words “repair of aneurysm” does not have a simple one-to-one correspondent in PCS. The I-9 description identifies the diagnosis of aneurysm, (information which should already be captured on the record in the diagnosis code) but does not actually give any indication of the specific method of repair. Depending on the documentation in the record, the correct PCS code could be one of several root operations: excision, replacement, or restriction, to name a few. All we know is that whatever is done to “repair” an aneurysm is included in the I-9 code description.

Procedure Codes and Levels of Specificity

I-9 and I-10 Code Sets Compared:
Code Length and Set Size

	ICD-9-CM	ICD-10-PCS
# of Characters	3-4 Numeric	7 Alphanumeric
# of Codes	~4,000	~73,000

As shown in the table above, PCS codes are longer, and there are many times more of them. Consequently, in an unabridged I-9 to PCS mapping, each I-9 code is typically linked to more than one PCS code, because each PCS code is more specific.

PCS is much more precise than I-9, and, just as important for purposes of mapping, the level of precision in a PCS code is standardized across the system. Within I-9, on the other hand, the level of detail varies greatly between codes. For example, category 39, Other operations on vessels, contains two codes with very different levels of detail:

- 39.31 Suture of artery
- 39.55 Reimplantation of aberrant renal vessel

The first code contains a precise description of the surgical technique (suture) but is very general with respect to location (an artery somewhere). The second code does not specify the method of reimplantation, but on other subjects is much more specific, containing a precise description of both the body part (renal vessel) and the diagnosis for which the procedure was performed (aberrant attachment to kidney, i.e., congenital anomaly).

I-9 descriptions or “includes” notes may contain several variations of a procedure. In practical terms this means that one general I-9 code actually represents a whole family of codes. Procedures that are identified by such “umbrella” codes lose their uniqueness as coded data. For example, an I-9 procedure code may include both the words “excision” and “destruction” of a body part in its description. This means that either an excision or a destruction procedure could have been performed. When only the coded I-9 data is available it is impossible to tell which method was used.

In PCS, by contrast, each significant variation is a unique procedure code and is defined consistently throughout the system. Excision and destruction are distinct root operations in PCS, identified as unique procedure codes.

One would not expect a PCS code to be linked to more than one I-9 code, since PCS is so much larger and more specific. However, since I-9 can be inconsistent, there are inevitable cases where it contains more detail than PCS. Characteristics of some I-9 procedure codes, such as subdividing body parts or approaches for a single procedure code category, or including diagnosis information in the description, were intentionally not incorporated in the PCS design. The ramification for mapping is that a PCS code may be linked to more than one I-9 code option, and choosing a closest match depends on the purpose of the mapping and whether the specific documentation in the chart is available.

Below are two examples where a distinction made in I-9 is not made in PCS. The result is that the PCS code may be linked to more than one I-9 code, because a particular area of the I-9 code set (highlighted in yellow) is more detailed than the norm.

Varying Specificity in I-9:
Body Part Subdivided

I-9 contains	I-9 also contains	PCS contains
83.64 Other suture of tendon	83.61 Suture of tendon sheath	0LQ70ZZ Repair Right Hand Tendon, Open Approach
85.23 Subtotal mastectomy	85.22 Resection of quadrant of breast	0HBT0ZZ Excision of Right Breast, Open Approach

Varying Specificity in I-9:
Approaches Unique to Pituitary Gland

I-9 contains	I-9 also contains	PCS contains
07.69 Total excision of pituitary gland, unspecified approach	07.64 Total excision of pituitary gland, transfrontal approach	0GT00ZZ Resection of Pituitary Gland, Open Approach
	07.65 Total excision of pituitary gland, transsphenoidal approach	
	07.68 Total excision of pituitary gland, other specified approach	

Procedure Codes and Approach

Approach is another area that complicates translating between I-9 and PCS. In PCS, approach is defined as “the technique used to reach the site of the procedure.” Further, all approaches used in PCS are defined, and these definitions aid in choosing the correct code. Two examples are “open” and “percutaneous.”

- *Open*—cutting through the skin or mucous membrane and any other body layers necessary to expose the site of the procedure
- *Percutaneous*—Entry, by puncture or minor incision, of instrumentation through the skin or mucous membrane and/or any other body layers necessary to reach the site of the procedure

By contrast, what constitutes “approach” is not defined in I-9, nor are the specific approaches used in I-9 codes defined (e.g., open, closed). Consequently the notion of approach itself is handled inconsistently in the system, and specific approaches can be difficult to interpret for correct coding.

For example, correct coding of an exploratory laparotomy followed by needle biopsy of the liver requires two separate codes in I-9:

54.11 Exploratory laparotomy
50.11 Closed (percutaneous) [needle] biopsy of liver

Though both codes appear to identify an operative approach, here they are not being used for the same purpose. The laparotomy code specifies the technique used to expose the site of the procedure. But the closed biopsy code specifies the instrument employed to obtain the biopsy, not the technique used to expose the procedure site. In other clinical situations, this same closed biopsy code specifies something different—a needle biopsy obtained through the skin without making an incision. Hence one I-9 code for “closed” biopsy of the liver can mean two very different things, depending on what other procedures were performed during the same operative episode.

In this example, correct coding in I-9 relies on a shifting notion of approach. By contrast, in PCS there is no ambiguity. If a biopsy of the liver is obtained by cutting through the skin and intervening tissue to expose the liver and then using a needle to take the sample, the approach selected for the PCS code is *open*. If a biopsy of the liver is obtained using a needle through the skin and intervening layers, the approach selected for the PCS code is *percutaneous*.

The comparative lack of precision between the two systems has ramifications for mapping. How one would map the procedure code for closed biopsy from I-9 to PCS depends on the wider clinical context of the operative episode. There is no simple “right” answer.

Procedure Codes in Combination: I-9 to PCS

Sometimes two procedures commonly performed together are identified in a single umbrella code, as in code 65.41, *Laparoscopic unilateral salpingo-oophorectomy*. I-9 also lists variations of combined procedures under a bigger umbrella code. This can be a general description, as in code 39.49, *Other revision of vascular procedure*, where varying combinations of multiple procedures could have been performed. Alternatively the procedure code may essentially say, “Here is the diagnosis, and this procedure code includes any of a number of things done to attempt to treat this condition.” It identifies the diagnosis clearly, but does not shed much light on the procedure. Examples include code 35.81, *Total repair of tetralogy of Fallot*, and code 03.53, *Repair of vertebral fracture*.

Mapping in the above cases, where according to the PCS definition of a procedure multiple procedures may be performed, requires that the I-9 code be linked to multiple PCS codes or ranges of codes. Because a PCS code identifies a single standardized classification of a single procedure, multiple PCS codes must be coded to fully describe the procedures performed in an I-9 combination code. Each PCS code is a partial expression of the information contained in the I-9 code. Entries of this type are linked using a special mapping attribute that indicates the allowable A+B+C choices.

In the table below, the same codes used in the above example, for laparoscopic salpingo-oophorectomy, are displayed with their full descriptions. Note that because the I-9 code includes procedures on two distinct body parts, two codes are required in PCS.

I-9 to PCS mapping:

“Laparoscopic salpingo-oophorectomy, bilateral”

ICD-9-CM Source	≈	ICD-10-PCS Target
65.63 Laparoscopic removal of both ovaries and tubes at same operative episode	≈	0UT24ZZ Resection of bilateral ovaries, percutaneous endoscopic approach AND 0UT74ZZ Resection of bilateral fallopian tubes, percutaneous endoscopic approach

Procedure Codes in Combination: PCS to I-9

A PCS code specifies a single procedure as defined within the PCS system. However, sometimes a PCS code must be linked to multiple I-9 codes because the I-9 primary procedure code is incomplete and so requires additional codes to convey specific information about the procedure. In I-9 these additional codes are referred to as “adjunct” procedure codes. They have been used increasingly in I-9, and function much like code extensions or modifiers in other systems: they convey additional information about the procedure performed, such as the number of devices placed or procedure sites treated. The detail contained in an I-9 primary procedure code plus an

I-9 adjunct code can be found in a single PCS code. The result is that one PCS code must sometimes be linked to a combination of I-9 codes—the principal procedure code plus adjunct code(s).

PCS to I-9 mapping:

“Percutaneous Transluminal Coronary Angioplasty (PTCA) of two coronary arteries, with insertion of two coronary stents”

ICD-10-PCS Source	≈	ICD-9-CM Target
02713DZ Dilation of coronary artery, two sites using intraluminal device, percutaneous approach	≈	00.66 PTCA or coronary atherectomy AND 00.41 Procedure on two vessels AND 00.46 Insertion of two vascular stents AND 36.06 Insertion of non-drug-eluting coronary artery stents

Introduction to the GEMs

The PCS and I-9 GEMs are used to facilitate linking between the procedure codes in I-9 volume 3 and the new PCS code set. The GEMs are formatted as downloadable “flat” text files. Each file contains a list of code pairs. Each code pair identifies a translation between a code in the source system and a code in the target system. First is the code in the source system, followed by a single code in the target system. The code in the source system is repeated on more than one line if there are multiple code alternatives in the target system, or if it is a combination entry. After the code pair come the attributes that apply to that code pair. The attributes can be used to analyze and reconcile the differences between the two coding systems. The GEMs are the raw material from which providers, health information vendors and payors can design and create specific mapping applications. This is covered in more detail in section 2.

The I-9 to I-10 GEM contains entries of interest if the coded medical record or test record generated for systems analysis uses I-9. The user could, for example, look at the GEM to see all possible options if the record had been coded in PCS, or to see if one I-9 test case will yield several alternative PCS test cases.

The PCS to I-9 GEM, on the other hand, contains entries of interest if the procedure coded, or the test case generated, uses PCS. In this case, the GEM facilitates mapping back to the I-9 alternatives that could represent how the PCS procedure might have been coded in I-9—for example, to determine how a procedure would be reimbursed.

The word “crosswalk” is often used to refer to mappings between annual code updates of I-9. Crosswalk carries with it a comfortable image: clean white lines mark the boundary on either side; the way across the street is the same in either direction; a traffic signal, or perhaps even a crossing guard, aids you from one side to the other. Please be advised: *GEMs are not crosswalks*. They are reference mappings, to help the user navigate the complexity of translating meaning from one code set to the other. They are tools to help the user understand, analyze, and make distinctions that manage the complexity, and to derive their own applied mappings if that is the goal. The GEMs are more complex than a simple one-to-one crosswalk, but ultimately more useful. They reflect the relative complexity of the code sets clearly so that it can be managed effectively, rather than masking it in an oversimplified way.

An entry in the GEM identifies relationships between a code in the source system and possible equivalents in the target system. If a mapping is described as having a direction, the source is the code one is mapping from, and the target is the code being mapped to.

	Source		Target		a.k.a.
From	ICD-9-CM	To	ICD-10-PCS		“forward mapping”
From	ICD-10-PCS	To	ICD-9-CM		“backward mapping”

The correspondence between codes in the source and target systems is approximate in most cases. As with translating between languages, translating between coding systems does not necessarily yield an exact match. Context is everything, and the specific purpose of an applied mapping must be identified before the most appropriate option can be selected.

The GEMs together provide a general (many to many) reference mapping that can be refined to fit the requirements of an applied mapping. For a particular code entry, the GEM may contain several possible translations, each on a separate row. The code in the source system is listed on a new row as many times as there are alternatives in the target system. Each translation is formatted as a code pair. The user must choose from among the alternatives a single code in the target system if a one-to-one mapping is desired.

The word “entry,” as used to describe the format of the GEM, refers to all rows in the GEM file having the same first listed code, the code in the source system. The word “row” refers to a single line in the file, containing a single code pair—one code from the source system and one code from the target system—along with its associated attributes. An entry typically encompasses multiple rows.

There are two basic types of entries in the GEM. They are “single entry” and “combination entry.” In special cases, a code in the source system may be mapped using both types of entries.

- *Single entry*—an entry in the GEM for which a code in the source system linked to one code option in the target system is a valid entry

An entry of the single type is characterized by a single translation: code A in the source system translates to code A **or** code B **or** code C in the target system. Each row in the entry can be one of several valid translations, and each is an option for a “one to one” applied mapping. An entry may consist of one row, if there is a close correspondence between the two codes in the code pair.

An entry of the single type is not the same as a one-to-one mapping. A code in the source system may be used multiple times in the GEM, each time linked to a different code in the target system. This is because the GEM contains alternative equivalent relationships from which the appropriate applied mapping can be selected. Taken together, all rows containing the same source system code linked to single code alternatives are considered one entry of the single type.

Here is an entry of the single type, consisting of three rows. The rows can be thought of as rows A **or** B **or** C. Each row of the entry is considered a valid applied mapping option.

I-9 to PCS GEM:

Single type entry for I-9 code 02.11

ICD-9-CM Source	≈	ICD-10-PCS Target
02.11 Simple suture of dura mater of brain	≈	00Q20ZZ Repair of dura mater, open approach
02.11 Simple suture of dura mater of brain	≈	00Q23ZZ Repair of dura mater, percutaneous approach
02.11 Simple suture of dura mater of brain	≈	00Q24ZZ Repair of dura mater, percutaneous endoscopic approach

Because PCS codes are for the most part more specific than I-9 codes, an entry of the single type in the I-9 to PCS GEM is typically linked to multiple PCS codes. The user must know, or must model, the level of detail contained in the original medical record to be able to settle on one of the PCS codes. The I-9 code itself cannot contain the answer; it cannot be made to describe detail it does not have. The same is occasionally true for the PCS to I-9 GEM as well. A PCS code may be linked to more than one I-9 code because of the variation in I-9 specificity and I-9’s use of adjunct codes, described in detail later.

Both I-9 and PCS contain what we refer to as “combination codes.” These are codes that contain more than one procedure in the code description. An example is PCS code *02733ZZ Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach*. In this case, I-9 does not have an equivalent combination code, so in order to link the PCS code to its I-9 equivalent, a combination entry must be used in the GEM.

- *Combination entry*—an entry in the GEM for which a code in the source system must be linked to more than one code option in the target system to be a valid entry

An entry of the combination type is characterized by a compound translation: code A in the source system must be linked simultaneously to code A **and** code B **and** code C in the target system to be a valid translation. Attributes in the GEM file clearly signal these special cases.

Stated another way, it takes more than one code in the target system to satisfy all of the meaning contained in one code in the source system. As discussed in this section, the situation occurs both when I-9 is the source system and when PCS is the source system.

Here is an entry of the combination type, consisting of two rows in the format of the GEM file. The rows can be thought of as rows A **and** B. The rows of the entry combined are considered one complete translation.

PCS to I-9 GEM:

Combination type entry for PCS code 02733ZZ

ICD-10-PCS Source	≈	ICD-9-CM Target
02733ZZ Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	≈	00.66 PTCA or coronary atherectomy AND 00.43 Procedure on four or more vessels

Linking a code in the source system to a combination of codes in the target system is accomplished by using conventions in the GEMs called *scenarios* and *choice lists*.

- *Scenario*—in a combination entry, a collection of codes from the target system containing the necessary codes that combined as directed will satisfy the equivalent meaning of a code in the source system

- *Choice list*—in a combination entry, a list of one or more codes in the target system from which one code must be chosen to satisfy the equivalent meaning of a code in the source system

Here is the combination type entry for PCS code *02733ZZ Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach* as it is depicted in the “flat text” GEM format, and repeated below in table format with the code descriptions and attributes labeled.

02733ZZ 0066 101 1 1
02733ZZ 0043 101 1 2

There are two rows in the PCS to I-9 GEM for combination code 02733ZZ. The entry is of the combination type, meaning that each row—code 02733ZZ linked to both of the two I-9 codes—is considered a valid entry. The combination flag is highlighted in the tables. The scenario number is 1, because there is only one version of the procedure specified in the combination code. There are two choice lists in this entry, and only one code in each choice list.

PCS Code	PCS Description	I-9 Code	I-9 Description	Approximate [FLAG]	No Map [FLAG]	Combination [FLAG]	Scenario	Choice list
02733ZZ	Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	00.66	PTCA or coronary atherectomy	1	0	1	1	1
02733ZZ	Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	00.43	Procedure on four or more vessels	1	0	1	1	2

It is important to make the distinction between a single row in a combination entry and an entry of the single type. An entry of the single type is one code in the source system linked to multiple one-code alternatives in the target system. It presents the option of linking one code in the source system to code **A or B or C** in the target system. Each code translation is considered a viable option. Each row of the source system code entry linked with target code **A or B or C** is one valid entry in an applied map.

An entry of the combination type is one code in the source system linked to a multiple-code alternative in the target system. If the source system is PCS, for example, the user **must** include I-9 codes **A and B and C** in order to cover all the diagnoses identified in the PCS code. Further, there may be more than one multiple-code alternative. If the GEM contains a range of I-9 code alternatives for each partial expression of the PCS code, then the number of solutions increases. Each instance of the PCS combination code paired with one code of the allowed range A and one code of the allowed range B and one code of the allowed range C is considered a valid entry. The combination flag in the GEM will clearly signal an entry of the combination type.

The two entry types and their main features are summarized in the table below.

Entry Type	Summary Description	Approximate [FLAG]	No Map [FLAG]	Combination [FLAG]	Scenario	Choice list
Single	Source system code has one or more single target code alternatives	On or Off	N/A	Off	0	0
Combination	Source system code has one or more multiple target code alternatives	On	N/A	On	1-9	1-9


Section 2—How to Use GEM Files

For ease of use, we recommend loading the GEM files into a database along with the code descriptions for both code sets. With roughly 90,000 codes and their descriptions in both code sets, a desktop database like MS Access is adequate.


ICD-10-PCS long format code descriptions can be found at:
http://www.cms.hhs.gov/ICD10/02_ICD-10-PCS.asp#TopOfPage

ICD-9-CM code descriptions can be found at:
http://www.cms.hhs.gov/ICD9ProviderDiagnosticCodes/06_codes.asp#TopOfPage

A general process for using the GEMs consists of three basic steps:



Step 1: EXTRACT



- *Have all rows that contain the same code from the source system been selected?*
- *Does the entry include multiple rows?*
- *Is the entry of the single type or combination type, or both?*

The code we will use for purposes of demonstration is I-9 code *02.11, Simple suture of dura mater of brain.*

I-9 to PCS GEM:

02.11 Simple suture of dura mater of brain

```


0207 0NP00JZ 10000
0207 0NP03JZ 10000
0207 0NP04JZ 10000
0211 00Q20ZZ 10000
0211 00Q23ZZ 10000
0211 00Q24ZZ 10000
0212 00Q10ZZ 10000
0212 00Q13ZZ 10000
0212 00Q14ZZ 10000
0212 00R107Z 10000
0212 00R10JZ 10000
0212 00R10KZ 10000
0212 00R147Z 10000
0212 00R14JZ 10000
0212 00R14KZ 10000
0212 00R207Z 10000
0212 00R20JZ 10000
0212 00R20KZ 10000
0212 00R247Z 10000
0212 00R24JZ 10000
0212 00R24KZ 10000
    
```

The illustrations at left and below display the I-9 procedure code 02.11 as it appears in the I-9 to PCS GEM. At left is the entry in text file format, and below is the same information as it would appear in a desktop database. Note that the codes do not contain decimals in the GEMs.

The code in the source system is listed first, followed by the code in the target system. Here the source system is the I-9 code and the target system is the PCS code. The final group of digits is used to indicate additional attributes for entries in the map. The first three digits are called flags. The last two digits are used in combination entries, and will be discussed later. This version of the GEM file contains a flag characterizing the degree of correspondence between codes in one row (“approximate” flag), a flag for codes with no translation in the target system (“no map” flag) and a flag indicating the row is part of a combination entry (“combination” flag). If the digit is 1, the flag applies (is “turned on”) to that entry in the GEM. If the digit is 0, the flag does not apply (is “turned off”) to that entry in the GEM. In other words, 1 means “yes,” the flag applies to the entry in the GEM and 0 means, “no,” the flag does not apply. There are three rows in I-9 to PCS GEM for code 02.11. The entry is of the single type, meaning that each row—code 02.11 linked to one of three PCS code alternatives—is considered a valid entry.

I-9 Code	I-9 Description	PCS Code	PCS Description	Approximate [FLAG]	No Map [FLAG]	Combination [FLAG]
02.11	Simple suture of dura mater of brain	00Q20ZZ	Repair of dura mater, open approach	1	0	0
02.11	Simple suture of dura mater of brain	00Q23ZZ	Repair of dura mater, percutaneous approach	1	0	0
02.11	Simple suture of dura mater of brain	00Q24ZZ	Repair of dura mater, percutaneous endoscopic approach	1	0	0

Step 2: ANALYZE



Is the “approximate” flag turned on?

- If yes, the translation is not a precise equivalent.

Is the “no map” flag turned on?

- If yes, there is no corresponding code in the target system.

Is the “combination” flag turned on?

- If yes, more than one code in the target system is required to satisfy the meaning of the code in the source system.

In the GEMs, there are three flags:

Approximate *indicates that the entry is not considered equivalent*

No Map *indicates that a code in the source system is not linked to any code in the target system*

Combination *indicates that more than one code in the target system is required to satisfy the full equivalent meaning of a code in the source system*

The Approximate Flag

I-9 Code	I-9 Description	PCS Code	PCS Description	Approximate [FLAG]	No Map [FLAG]	Combination [FLAG]
02.11	Simple suture of dura mater of brain	00Q20ZZ	Repair of dura mater, open approach	1	0	0
02.11	Simple suture of dura mater of brain	00Q23ZZ	Repair of dura mater, percutaneous approach	1	0	0
02.11	Simple suture of dura mater of brain	00Q24ZZ	Repair of dura mater, percutaneous endoscopic approach	1	0	0

The approximate flag is turned on when no one code in the target system or linked combination of codes in the target system expresses the same essential meaning as the code in the source system. Because the I-9 and PCS structure and organization are so different, this flag is turned on for the great majority of entries in the GEMs. The difference between the two systems is typically in level of detail between the codes, and in nearly all cases the PCS code is more detailed than the I-9 code. The approximate flag is turned on for all rows in the GEM for I-9

code 02.11. The level of detail differs here—the approach (highlighted in the table) is specified in PCS and not in I-9.

The No Map Flag

I-9 Code	I-9 Description	PCS Code	PCS Description	Approximate [FLAG]	No Map [FLAG]	Combination [FLAG]
00.40	Procedure on single vessel	NoPCS	No Description	0	1	0
00.41	Procedure on two vessels	NoPCS	No Description	0	1	0
00.42	Procedure on three vessels	NoPCS	No Description	0	1	0
00.43	Procedure on four or more vessels	NoPCS	No Description	0	1	0
00.44	Procedure on vessel bifurcation	NoPCS	No Description	0	1	0

In the I-9 to PCS GEM, the “no map” flag is on for a subset of I-9 codes. I-9 contains adjunct codes that do not identify a procedure, but instead further specify an aspect of a procedure, such as the number of stents used in an angioplasty. They must be paired with an I-9 “primary procedure” code to be meaningful. Since they cannot be coded alone in I-9, they cannot be linked singly to an equivalent code in PCS, because every PCS code is complete in itself as a procedure code. To put it another way, there are no PCS adjunct codes; every PCS code is a primary procedure code. In the I-9 to PCS GEM, I-9 adjunct codes are listed without a corresponding PCS entry, and with the “no map” flag on.

The adjunct I-9 codes are listed in the table below. In the I-9 to PCS GEM, the “no map” flag is on for adjunct I-9 codes and they are not linked to any PCS code. In the PCS to I-9 GEM, adjunct I-9 codes are included as linked portions of an I-9 combination entry.

Code	Adjunct ICD-9-CM codes Description
00.40	Procedure on single vessel
00.41	Procedure on two vessels
00.42	Procedure on three vessels
00.43	Procedure on four or more vessels
00.44	Procedure on vessel bifurcation
00.45	Insertion of one vascular stent
00.46	Insertion of two vascular stents
00.47	Insertion of three vascular stents
00.48	Insertion of four or more vascular stents
00.55	Insertion of drug-eluting peripheral vessel stent(s)
00.63	Percutaneous insertion of carotid artery stent(s)
00.64	Percutaneous insertion of other precerebral (extracranial) artery stent(s)
00.65	Percutaneous insertion of intracranial vascular stent(s)
00.74	Hip replacement bearing surface, metal on polyethylene
00.75	Hip replacement bearing surface, metal-on-metal
00.76	Hip replacement bearing surface, ceramic-on-ceramic
00.77	Hip replacement bearing surface, ceramic-on- polyethylene
00.91	Transplant from live related donor
00.92	Transplant from live non-related donor
00.93	Transplant from cadaver
36.06	Insertion of non-drug-eluting coronary artery stent(s)
36.07	Insertion of drug-eluting coronary artery stent(s)
39.90	Insertion of non-drug-eluting peripheral vessel stent(s)
70.94	Insertion of biological graft
70.95	Insertion of synthetic graft or prosthesis
72.4	Forceps rotation of fetal head
81.62	Fusion or refusion of 2-3 vertebrae
81.63	Fusion or refusion of 4-8 vertebrae
81.64	Fusion or refusion of 9 or more vertebrae
84.51	Insertion of interbody spinal fusion device

The Combination Flag

The combination flag is turned on when a code in the source system must be linked to more than one code in the target system to be a valid entry. When the combination flag is on, the *scenario* and *choice list* fields in the GEM contain a number. They appear last in the GEM text file, after the flags. These numbers allow the user to collate the combination entries in the GEM.

```
304 OCTS0ZZ 10111
304 OCTS4ZZ 10111
304 OCTS7ZZ 10111
304 OCTS8ZZ 10111
304 0WB60ZZ 10112
304 0WB63ZZ 10112
304 0WB64ZZ 10112
304 0WB6XZZ 10112
304 OCTS0ZZ 10121
304 OCTS4ZZ 10121
304 OCTS7ZZ 10121
304 OCTS8ZZ 10121
304 0WB60ZZ 10122
304 0WB63ZZ 10122
304 0WB64ZZ 10122
304 0WB6XZZ 10122
304 0GTG0ZZ 10123
304 0GTG4ZZ 10123
304 0GTH0ZZ 10123
304 0GTH4ZZ 10123
304 0GTK0ZZ 10123
304 0GTK4ZZ 10123
```

I-9 to PCS GEM:

30.4 Radical Laryngectomy

Complete [total] laryngectomy with radical neck dissection (with thyroidectomy)
(with synchronous tracheostomy)

The illustrations at left and below display the entry for I-9 procedure code 30.4, *Radical Laryngectomy*, as it appears in the I-9 to PCS GEM. At left is the entry in text file format, and below is the same information as it would appear in a desktop database. The I-9 procedure code 30.4 describes more than one procedure in PCS, so it requires a combination entry in the GEM. A combination is subdivided hierarchically on two levels: 1) By *scenario*, the number of variations of procedure combinations included in the source system code, and 2) By *choice list*, the possible target system codes that combined are one valid expression of a scenario. Each procedure listed in the “includes” notes of the I-9 code is a unique PCS code, so more than one PCS code is required to satisfy the equivalent meaning. Therefore, each PCS code for a different procedure is assigned a *choice list* number in the GEM.

In addition, three distinct clinical variations of the procedure are specified in I-9 code 30.4: one includes the laryngectomy and neck dissection, the second adds resection of the thyroid, and the third adds a tracheostomy. The linking between I-9 code 30.4 and PCS that includes these additional procedures must be distinguished from the linking that does not. Each clinically distinct variation of a procedure combination requires its own corresponding list of codes in PCS, so each version of the procedure is assigned a separate *scenario* number in the GEM.

A scenario designates one version of all the source system procedures as specified in a combination code. In other words, it identifies one roughly equivalent expression of the source system code. In this example, scenario 1 contains all the PCS codes needed to satisfy the equivalent meaning of “Complete laryngectomy with radical neck dissection.” Scenario 2 contains all the PCS codes needed for “Complete laryngectomy with radical neck dissection with thyroidectomy.” Scenario 3 contains all the PCS codes needed for “Complete laryngectomy with radical neck dissection with synchronous tracheostomy.”

A scenario is subdivided into two or more choice lists of codes in the target system. These are the codes that must be linked together in an applied mapping to satisfy the equivalent meaning of the code in the source system. A choice list contains one or more codes in the target system that

express a portion of the meaning of the code in the source system. A code must be included from each choice list in a scenario to satisfy the equivalent meaning of the code in the source system.

Scenario 1

30.4 Radical Laryngectomy

Complete [total] laryngectomy with radical neck dissection

I-9 Code	I-9 Description	PCS Code	PCS Description	Approximate [FLAG]	No Map [FLAG]	Combination [FLAG]	Scenario	Choice list
30.4	Radical laryngectomy	0CTS0ZZ	Resection of Larynx, Open Approach	1	0	1	1	1
30.4	Radical laryngectomy	0CTS4ZZ	Resection of Larynx, Percutaneous Endoscopic Approach	1	0	1	1	1
30.4	Radical laryngectomy	0CTS7ZZ	Resection of Larynx, Via Natural or Artificial Opening	1	0	1	1	1
30.4	Radical laryngectomy	0CTS8ZZ	Resection of Larynx, Via Natural or Artificial Opening Endoscopic	1	0	1	1	1
30.4	Radical laryngectomy	0WB60ZZ	Excision of Neck, Open Approach	1	0	1	1	2
30.4	Radical laryngectomy	0WB63ZZ	Excision of Neck, Percutaneous Approach	1	0	1	1	2
30.4	Radical laryngectomy	0WB64ZZ	Excision of Neck, Percutaneous Endoscopic Approach	1	0	1	1	2

In this example there are two PCS choice lists in scenario 1, three PCS choice lists in scenario 2, and three PCS choice lists in scenario 3.

Scenario 2

30.4 Radical Laryngectomy

Complete [total] laryngectomy with radical neck dissection with thyroidectomy

I-9 Code	I-9 Description	PCS Code	PCS Description	Approximate [FLAG]	No Map [FLAG]	Combination [FLAG]	Scenario	Choice list
30.4	Radical laryngectomy	0CTS0ZZ	Resection of Larynx, Open Approach	1	0	1	2	1
30.4	Radical laryngectomy	0CTS4ZZ	Resection of Larynx, Percutaneous Endoscopic Approach	1	0	1	2	1
30.4	Radical laryngectomy	0CTS7ZZ	Resection of Larynx, Via Natural or Artificial Opening	1	0	1	2	1
30.4	Radical laryngectomy	0CTS8ZZ	Resection of Larynx, Via Natural or Artificial Opening Endoscopic	1	0	1	2	1
30.4	Radical laryngectomy	0WB60ZZ	Excision of Neck, Open Approach	1	0	1	2	2
30.4	Radical laryngectomy	0WB63ZZ	Excision of Neck, Percutaneous Approach	1	0	1	2	2
30.4	Radical laryngectomy	0WB64ZZ	Excision of Neck, Percutaneous Endoscopic Approach	1	0	1	2	2
30.4	Radical laryngectomy	0WB6XZZ	Excision of Neck, External Approach	1	0	1	2	2
30.4	Radical laryngectomy	0GTG0ZZ	Resection of Left Thyroid Gland Lobe, Open Approach	1	0	1	2	3
30.4	Radical laryngectomy	0GTG4ZZ	Resection of Left Thyroid Gland Lobe, Percutaneous Endoscopic Approach	1	0	1	2	3
30.4	Radical laryngectomy	0GTH0ZZ	Resection of Right Thyroid Gland Lobe, Open Approach	1	0	1	2	3
30.4	Radical laryngectomy	0GTH4ZZ	Resection of Right Thyroid Gland Lobe, Percutaneous Endoscopic Approach	1	0	1	2	3
30.4	Radical laryngectomy	0GTK0ZZ	Resection of Thyroid Gland, Open Approach	1	0	1	2	3
30.4	Radical laryngectomy	0GTK4ZZ	Resection of Thyroid Gland, Percutaneous Endoscopic Approach	1	0	1	2	3

Scenario 3

30.4 Radical Laryngectomy

Complete [total] laryngectomy with radical neck dissection with synchronous tracheostomy

I-9 Code	I-9 Description	PCS Code	PCS Description	Approximate [FLAG]	No Map [FLAG]	Combination [FLAG]	Scenario	Choice list
30.4	Radical laryngectomy	0CTS0ZZ	Resection of Larynx, Open Approach	1	0	1	3	1
30.4	Radical laryngectomy	0CTS4ZZ	Resection of Larynx, Percutaneous Endoscopic Approach	1	0	1	3	1
30.4	Radical laryngectomy	0CTS7ZZ	Resection of Larynx, Via Natural or Artificial Opening	1	0	1	3	1
30.4	Radical laryngectomy	0CTS8ZZ	Resection of Larynx, Via Natural or Artificial Opening Endoscopic	1	0	1	3	1
30.4	Radical laryngectomy	0WB60ZZ	Excision of Neck, Open Approach	1	0	1	3	2
30.4	Radical laryngectomy	0WB63ZZ	Excision of Neck, Percutaneous Approach	1	0	1	3	2
30.4	Radical laryngectomy	0WB64ZZ	Excision of Neck, Percutaneous Endoscopic Approach	1	0	1	3	2
30.4	Radical laryngectomy	0WB6XZZ	Excision of Neck, External Approach	1	0	1	3	2
30.4	Radical laryngectomy	0B110F4	Bypass Trachea to Cutaneous with Tracheostomy Device, Open Approach	1	0	1	3	3
30.4	Radical laryngectomy	0B110Z4	Bypass Trachea to Cutaneous, Open Approach	1	0	1	3	3
30.4	Radical laryngectomy	0B113F4	Bypass Trachea to Cutaneous with Tracheostomy Device, Percutaneous Approach	1	0	1	3	3
30.4	Radical laryngectomy	0B113Z4	Bypass Trachea to Cutaneous, Percutaneous Approach	1	0	1	3	3
30.4	Radical laryngectomy	0B114F4	Bypass Trachea to Cutaneous with Tracheostomy Device, Percutaneous Endoscopic Approach	1	0	1	3	3
30.4	Radical laryngectomy	0B114Z4	Bypass Trachea to Cutaneous, Percutaneous Endoscopic Approach	1	0	1	3	3

Step 3: REFINE

REFINE
Select the row(s) of an entry that meet the requirements of an applied mapping.

- *What is the purpose of the applied mapping?*
- *Does the applied mapping require that the code in the source system be mapped to only one “best” alternative in the target system?*
- *Will the correct applied mapping vary depending on the documentation in the record?*

Once the user has analyzed all rows for an entry in the GEM, it is possible to select the row or rows most appropriate to a specific mapping application. We will use two different sample entries of the combination type—one from the I-9 to PCS GEM and one from the PCS to I-9 GEM—in order to discuss the process of refining an entry and deriving an applied mapping.

Sample Entry 1—I-9 to PCS GEM

00.53 *Implantation or replacement of CRT pacemaker generator*

I-9 Code	I-9 Description	PCS Code	PCS Description	Approximate [FLAG]	No Map [FLAG]	Combination [FLAG]	Scenario	Choice list
00.53	Implantation <u>or</u> replacement of CRT pacemaker generator	0JH60P3	Insertion of Cardiac Resynchronization Pacemaker Pulse Generator into Chest Subcutaneous Tissue and Fascia, Open Approach	1	0	0	0	0
00.53	Implantation <u>or</u> replacement of CRT pacemaker generator	0JH63P3	Insertion of Cardiac Resynchronization Pacemaker Pulse Generator into Chest Subcutaneous Tissue and Fascia, Percutaneous Approach	1	0	0	0	0
00.53	Implantation <u>or</u> replacement of CRT pacemaker generator	0JPT0PZ	Removal of Pacemaker / Defibrillator from Trunk Subcutaneous Tissue and Fascia, Open Approach	1	0	1	1	1
00.53	Implantation <u>or</u> replacement of CRT pacemaker generator	0JPT3PZ	Removal of Pacemaker / Defibrillator from Trunk Subcutaneous Tissue and Fascia, Percutaneous Approach	1	0	1	1	1
00.53	Implantation <u>or</u> replacement of CRT pacemaker generator	0JH60P3	Insertion of Cardiac Resynchronization Pacemaker Pulse Generator into Chest Subcutaneous Tissue and Fascia, Open Approach	1	0	1	1	2
00.53	Implantation <u>or</u> replacement of CRT pacemaker generator	0JH63P3	Insertion of Cardiac Resynchronization Pacemaker Pulse Generator into Chest Subcutaneous Tissue and Fascia, Percutaneous Approach	1	0	1	1	2

In this instance an I-9 entry in the GEM is of both the single and the combination type. Because the I-9 code includes either implantation of the CRT pacemaker generator alone or removal of the old generator and insertion of a new one in the same operative episode, the GEM must translate both possibilities into their PCS equivalents. One version requires one PCS code (“implantation” in the I-9 code) to satisfy the equivalent meaning, and the other version requires two PCS codes (“replacement” in the I-9 code).

After classifying the entry into its single and combination entry constituents and collating the combination entries into their respective choice lists (there is only one combination scenario here), the available possibilities are:

Single entry

ICD-9-CM Source	≈	ICD-10-PCS Target
00.53 Implantation <u>or</u> replacement of CRT pacemaker generator	≈	0JH60P3 Insertion of Cardiac Resynchronization Pacemaker Pulse Generator into Chest Subcutaneous Tissue and Fascia, Open Approach OR 0JH63P3 Insertion of Cardiac Resynchronization Pacemaker Pulse Generator into Chest Subcutaneous Tissue and Fascia, Percutaneous Approach

OR

Combination entry

ICD-9-CM Source	≈	ICD-10-PCS Target
00.53 Implantation <u>or</u> replacement of CRT pacemaker generator	≈	0JPT0PZ Removal of Pacemaker / Defibrillator from Trunk Subcutaneous Tissue and Fascia, Open Approach OR 0JPT3PZ Removal of Pacemaker / Defibrillator from Trunk Subcutaneous Tissue and Fascia, Percutaneous Approach AND 0JH60P3 Insertion of Cardiac Resynchronization Pacemaker Pulse Generator into Chest Subcutaneous Tissue and Fascia, Open Approach OR 0JH63P3 Insertion of Cardiac Resynchronization Pacemaker Pulse Generator into Chest Subcutaneous Tissue and Fascia, Percutaneous Approach

Note that either member of choice list 1 can be combined with either member of choice list 2. Although PCS codes having the same approach value (5th character identifies operative approach for all Medical and Surgical and related codes) are more likely to be used together, the GEM must accommodate all possible variations of an operative episode and allow for combinations

with differing approach values. In this case a variation could be that the old generator was removed percutaneously and the new one inserted using an open approach.

To refine this entry, first the user must decide whether or not the applied mapping is going to encompass both the single and combination translation. This decision of course depends on the mapping application.

A health information professional who is converting old I-9 records to PCS, and has access to the individual medical record, can make use of the increased specificity in PCS codes. The user can simply refer to the original record to see the specific nature of the procedure and assign the correct PCS code(s) to the record.

However, a health statistics analyst or data modeler who is translating aggregate I-9 data forward to PCS, and has no access to individual medical records, cannot make use of the fine distinctions in PCS, since they are not present in the old data. In this case, finding the closest equivalent cannot be the goal. The user must choose a PCS code or pair of codes to represent all the possibilities, and could choose to fashion a rule by which to map similar cases. Rules specific to the applied mapping would promote consistency and document the decisions made. For example, here the applied mapping could use only the PCS Insertion codes.

A reimbursement specialist looking to translate a PCS coded record back to I-9 codes to produce equivalent payment from a legacy payment system does not need to retain the fine distinctions in PCS, since they are not present in the old system and will not affect payment. This user may want to ignore the recommended combinations and choose one row of the entry to link to the I-9 code that will represent all the PCS possibilities. In this case finding the one closest equivalent is not the goal.

However, a reimbursement specialist looking to upgrade a legacy payment system to make it more accurate and fair will want to use the increased specificity available in the PCS system. This user may be assigning new payment adjudication logic to the system (payment for insertion of the pacemaker could be reimbursed at a lower rate than removal of the old pacemaker and insertion of a new one). In this case, the I-9 to PCS GEM would be used as a reference, to see the PCS possibilities and assign them accordingly in a new reimbursement system, where records would be coded in PCS and paid using PCS codes.

Sample Entry 2—PCS to I-9 GEM:

02733ZZ Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach

PCS Code	PCS Description	I-9 Code	I-9 Description	Approximate [FLAG]	Combination [FLAG]	Scenario	Choice list
02733ZZ	Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	00.66	Percutaneous transluminal coronary angioplasty [PTCA] or coronary atherectomy	1	1	1	1
02733ZZ	Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	00.40	Procedure on single vessel	1	1	1	2
02733ZZ	Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	00.41	Procedure on two vessels	1	1	1	2
02733ZZ	Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	00.42	Procedure on three vessels	1	1	1	2
02733ZZ	Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	00.43	Procedure on four or more vessels	1	1	1	2
02733ZZ	Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	00.44	Procedure on vessel bifurcation	1	1	1	2

The approximate flag is on, indicating that the relationship between the code in the source system and the code in the target system is classified as an approximate equivalent only. In this case the difference lies in the classification of the body part. I-9 classifies the body part by number of **vessels** treated, and PCS classifies by the number of **sites** treated regardless of the number of vessels. For example, a PTCA could treat two separate lesions along the same vessel. In PCS this is considered two sites for coding purposes, whereas in I-9 it is considered one vessel. The ramification for mapping is that the PCS code indicating that four sites were treated must be linked to all I-9 code alternatives indicating the number of vessels treated, since all four sites could conceivably be on the same vessel.

The combination flag is on, and the number in the scenario column indicates there is only one variation of the procedure identified in the source system. After collating the rows of this entry into their respective choice lists, then, the available possibilities are:

ICD-10-PCS Source	≈	ICD-9-CM Target
02733ZZ Dilation of Coronary Artery, Four or More Sites, Percutaneous Approach	≈	00.66 PTCA or coronary atherectomy AND 00.40 Procedure on single vessel OR 00.41 Procedure on two vessels OR 00.42 Procedure on three vessels OR 00.43 Procedure on four or more vessels OR 00.44 Procedure on vessel bifurcation

There is only one way to refine this entry: to decide among the alternatives in choice list 2, the user must have access to the detail in the original record. If the patient record is not available, or the mapping application is intended to establish general rules for translation rather than deciding on a case-by-case basis, then a consistent method must be derived and documented for resolving the disparity in body part classification between the two systems. Depending on the mapping application, the user may want to equate vessels with sites or ignore the I-9 adjunct codes altogether in the applied mapping. These decisions require considering the ramifications of lost detail for accurate reimbursement—for coding patient records in one system when the bill will be submitted in another system—and for transposing research data gathered in one system and converted to another.

Glossary

Approach—a character of the seven-character ICD-10-PCS code that “defines the technique used to reach the site of the procedure”

Approximate flag—attribute in a GEM that when turned on indicates that the entry is not considered equivalent

Applied mapping—distillation of a reference mapping to conform to the needs of a particular application (i.e., data quality, reimbursement, research)

Backward mapping—mapping that proceeds from a newer code set to an older code set

Choice list—in a combination entry, a list of one or more codes in the target system from which one code must be chosen to satisfy the equivalent meaning of a code in the source system

Combination flag—attribute in a GEM that when turned on indicates that more than one code in the target system is required to satisfy the full equivalent meaning of a code in the source system

Combination entry—an entry in the GEM for which a code in the source system must be linked to more than one code option in the target system to be a valid entry

Forward mapping—mapping that proceeds from an older code set to a newer code set

General Equivalence Mapping (GEM)—reference mapping that attempts to include all valid relationships between the codes in the ICD-9-CM procedure classification and the ICD-10 Procedure Code System (ICD-10-PCS)

ICD-9-CM—International Classification of Diseases 9th Revision Clinical Modification (I-9)

ICD-10-PCS—ICD-10 Procedure Code System (PCS)

No map flag—attribute in a GEM that when turned on indicates that a code in the source system is not linked to any code in the target system

Reference mapping—mapping that includes all possible valid relationships between a source system and a target system

Root operation—a character of the seven-character ICD-10-PCS code that “defines the objective of the procedure”

Scenario—in a combination entry, a collection of codes from the target system containing the necessary codes that when combined as directed will satisfy the equivalent meaning of a code in the source system

Single entry—an entry in the GEM for which a code in the source system linked to one code option in the target system is a valid entry

Source system—code set of origin in the mapping; the set being mapped ‘from’

Target system—destination code set in the mapping; the set being mapped ‘to’

Procedure Code Set General Equivalence Mappings 2009 Version Documentation

Appendix A—File and Format Detail

ICD-9-CM to ICD-10-PCS General Equivalence Map (GEM) FILE AND FORMAT

FILE NAME: gem_i9pcs.txt

FILE FORMAT:

FIELD	POSITION	LENGTH	VALUE
ICD-9-CM Code [source]	1 – 5	5	Left justified, blank filled No decimal
<i>Filler</i>	6	1	<i>Blank</i>
ICD-10-PCS Code [target]	7 – 13	7	All seven characters used
<i>Filler</i>	14	1	<i>Blank</i>
Approximate [FLAG]	15	1	1 = Yes/On 0 = No/Off
No Map [FLAG]	16	1	1 = Yes/On 0 = No/Off
Combination [FLAG]	17	1	1 = Yes/On 0 = No/Off
Scenario	18	1	0 – 9
Choice list	19	1	0 – 9

**ICD-10-PCS to ICD-9-CM
General Equivalence Map (GEM)
FILE AND FORMAT**

FILE NAME: gem_pcsi9.txt

FILE FORMAT:

FIELD	POSITION	LENGTH	VALUE
ICD-10-PCS Code [source]	1 – 7	7	Left justified, blank filled No decimal
<i>Filler</i>	8	1	<i>Blank</i>
ICD-9-CM Code [target]	9 – 13	5	All seven characters used
<i>Filler</i>	14	1	<i>Blank</i>
Approximate [FLAG]	15	1	1 = Yes/On 0 = No/Off
No Map [FLAG]	16	1	1 = Yes/On 0 = No/Off
Combination [FLAG]	17	1	1 = Yes/On 0 = No/Off
Scenario	18	1	0 – 9
Choice list	19	1	0 – 9

Appendix B

New ICD-9-CM Code Entries in the ICD-9-CM to ICD-10-PCS GEM

This table contains the new I-9 codes valid on October 1, 2008, and their corresponding entries in the I-9 to PCS GEM. In most cases, the I-9 code is linked to more than one PCS code. The number of PCS codes in an entry is listed in the third column of the table if the codes are new for 2009, or in the fourth column if the codes are not new.

Where the GEM entry consists of one or two PCS codes, the code(s) and their descriptions are given in full. Where the GEM entry consists of more than two PCS codes, the applicable PCS codes are given in condensed notation (e.g. 02HW[0,3-4]2Z). To derive all valid codes in the entry, combine each character in each set of brackets with the non-bracketed characters in the correct order. The description is given for the first PCS code in the mapping entry, followed by the number of other codes contained in the entry (e.g., Insertion of Monitoring Device into Thoracic Aorta, Open Approach + 2 others). The values underlined in the description correspond to the first value in each bracket.

New ICD-9-CM	ICD-10-PCS	# New PCS Codes	# PCS Codes	New PCS Value
00.49 SuperSaturated oxygen therapy	5A0512C Extracorporeal Supersaturated Oxygenation, Intermittent; 5A0522C Extracorporeal Supersaturated Oxygenation, Continuous	2		[Char 7] C Supersaturated
00.58 Insertion of intra-aneurysm sac pressure monitoring device (intraoperative)	02HW[0,3-4]2Z Insertion of Monitoring Device into Thoracic Aorta, <u>Open</u> Approach + 2 others; 04H0[0,3-4]2Z Insertion of Monitoring Device into Abominal Aorta, <u>Open</u> Approach + 2 others;		6	
00.59 Intravascular pressure measurement of coronary arteries	4A03[0,3]BC Measurement of Arterial Pressure, Coronary, <u>Open</u> Approach + 1 other; 4A13[0,3]BC Monitoring of Arterial Pressure, Coronary, <u>Open</u> Approach + 1 other		4	
00.67 Intravascular pressure measurement of intrathoracic arteries	4A030BF Measurement of Arterial Pressure, Other Thoracic, Open Approach; 4A033BF Measurement of Arterial Pressure, Other Thoracic, Percutaneous Approach	2		[Char 7] F Other Thoracic
00.68 Intravascular pressure measurement of peripheral arteries	4A030B1 Measurement of Arterial Pressure, Peripheral, Open Approach; 4A033B1 Measurement of Arterial Pressure, Peripheral, Percutaneous Approach		2	
00.69 Intravascular pressure measurement, other specified and unspecified vessels	4A040B1 Measurement of Venous Pressure, <u>Peripheral</u> , <u>Open</u> Approach + 3 others; 4A140B1 Monitoring of Venous Pressure, <u>Peripheral</u> , <u>Open</u> Approach + 3 others		8	
17.11 Laparoscopic repair of direct inguinal hernia with graft or prosthesis	0YU[5,6]4[7,J-K]Z Supplement <u>Right</u> Inguinal Region with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 5 others		6	
17.12 Laparoscopic repair of indirect inguinal hernia with graft or prosthesis	0YU[5,6]4[7,J-K]Z Supplement <u>Right</u> Inguinal Region with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 5 others		6	
17.13 Laparoscopic repair of inguinal hernia with graft or prosthesis, not otherwise specified	0YU[5,6]4[7,J-K]Z Supplement <u>Right</u> Inguinal Region with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 5 others		6	

New ICD-9-CM	ICD-10-PCS	# New PCS Codes	# PCS Codes	New PCS Value
17.21 Laparoscopic bilateral repair of direct inguinal hernia with graft or prosthesis	0YUA4[7,J-K]Z Supplement Bilateral Inguinal Region with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 2 others		3	
17.22 Laparoscopic bilateral repair of indirect inguinal hernia with graft or prosthesis	0YUA4[7,J-K]Z Supplement Bilateral Inguinal Region with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 2 others		3	
17.23 Laparoscopic bilateral repair of inguinal hernia, one direct and one indirect, with graft or prosthesis	0YUA4[7,J-K]Z Supplement Bilateral Inguinal Region with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 2 others		3	
17.24 Laparoscopic bilateral repair of inguinal hernia with graft or prosthesis, not otherwise specified	0YUA4[7,J-K]Z Supplement Bilateral Inguinal Region with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 2 others		3	
17.31 Laparoscopic multiple segmental resection of large intestine	0DB[E-G]4ZZ Excision of <u>Large Intestine</u> , Percutaneous Endoscopic Approach + 2 others		3	
17.32 Laparoscopic cecectomy	0DTH4ZZ Resection of Cecum, Percutaneous Endoscopic Approach		1	
17.33 Laparoscopic right hemicolectomy	0DTF4ZZ Resection of Right Large Intestine, Percutaneous Endoscopic Approach		1	
17.34 Laparoscopic resection of transverse colon	0DTL4ZZ Resection of Transverse Colon, Percutaneous Endoscopic Approach		1	
17.35 Laparoscopic left hemicolectomy	0DTG4ZZ Resection of Left Large Intestine, Percutaneous Endoscopic Approach		1	
17.36 Laparoscopic sigmoidectomy	0DTN4ZZ Resection of Sigmoid Colon, Percutaneous Endoscopic Approach		1	
17.39 Other laparoscopic partial excision of large intestine	0DB[E-H,K-N]4ZZ; 0DT[K,M]4ZZ Excision of <u>Large Intestine</u> , Percutaneous Endoscopic Approach + 9 others		10	
17.41 Open robotic assisted procedure	8E0[9,W-Y]0CZ Robotic Assisted Procedure of <u>Head and Neck Region</u> , Open Approach + 3 others	4		[Char 7] C Robotic Assisted Procedure
17.42 Laparoscopic robotic assisted procedure	8E0W4CZ Robotic Assisted Procedure of Trunk Region, Percutaneous Endoscopic Approach	1		[Char 7] C Robotic Assisted Procedure
17.43 Percutaneous robotic assisted procedure	8E0[9,W-Y]3CZ Robotic Assisted Procedure of <u>Head and Neck Region</u> , Percutaneous Approach + 3 others	4		[Char 7] C Robotic Assisted Procedure
17.44 Endoscopic robotic assisted procedure	8E0[9,X-Y]4CZ Robotic Assisted Procedure of <u>Head and Neck Region</u> , Percutaneous Endoscopic Approach + 2 others; 8E0[9,W]8CZ Robotic Assisted Procedure of <u>Head and Neck Region</u> , Via Natural or Artificial Opening Endoscopic + 1 other	5		[Char 7] C Robotic Assisted Procedure
17.45 Thoracoscopic robotic assisted procedure	8E0W4CZ Robotic Assisted Procedure of Trunk Region, Percutaneous Endoscopic Approach	1		[Char 7] C Robotic Assisted Procedure

New ICD-9-CM	ICD-10-PCS	# New PCS Codes	# PCS Codes	New PCS Value
17.49 Other and unspecified robotic assisted procedure	8E0[9,W-Y]XCZ Robotic Assisted Procedure of Head and Neck Region + 3 others; 8E0[9,W]7CZ Robotic Assisted Procedure of <u>Head and Neck Region, Via Natural or Artificial Opening</u> + 1 other	6		[Char 7] C Robotic Assisted Procedure
33.72 Endoscopic pulmonary airway flow measurement	4A0985Z Measurement of Respiratory Flow, Via Natural or Artificial Opening Endoscopic		1	
37.36 Excision or destruction of left atrial appendage (LAA)	02B73ZZ Excision of Left Atrium, Percutaneous Approach		1	
37.55 Removal of internal biventricular heart replacement system	02PA[0,3-4]QZ Removal of Implantable Heart Assist System from Heart, <u>Open Approach</u> + 2 others		3	
37.60 Implantation or insertion of biventricular external heart assist system	02HA[0,3-4]RS Insertion of Biventricular External Heart Assist System into Heart, Open + 2 others	3		[Char 7] S Biventricular
38.23 Intravascular spectroscopy	8E023DZ Near Infrared Spectroscopy of Circulatory System, Percutaneous Approach	1		[Char 6] D Near Infrared Spectroscopy
45.81 Laparoscopic total intra-abdominal colectomy	0DTE4ZZ Resection of Large Intestine, Percutaneous Endoscopic Approach		1	
45.82 Open total intra-abdominal colectomy	0DTE0ZZ Resection of Large Intestine, Open Approach		1	
45.83 Other and unspecified total intraabdominal colectomy	0DTE7ZZ Resection of Large Intestine, Via Natural or Artificial Opening; 0DTE8ZZ Resection of Large Intestine, Via Natural or Artificial Opening Endoscopic		2	
48.40 Pull-through resection of rectum, not otherwise specified	0DTP0ZZ Resection of Rectum, Open Approach; 0DTP4ZZ Resection of Rectum, Percutaneous Endoscopic Approach		2	
48.42 Laparoscopic pull-through resection of rectum	0DTP4ZZ Resection of Rectum, Percutaneous Endoscopic Approach		1	
48.43 Open pull-through resection of rectum	0DTP0ZZ Resection of Rectum, Open Approach		1	
48.50 Abdominoperineal resection of the rectum, not otherwise specified	0DTP0ZZ Resection of Rectum, Open Approach; 0DTP4ZZ Resection of Rectum, Percutaneous Endoscopic Approach		2	
48.51 Laparoscopic abdominoperineal resection of the rectum	0DTP4ZZ Resection of Rectum, Percutaneous Endoscopic Approach		1	
48.52 Open abdominoperineal resection of the rectum	0DTP0ZZ Resection of Rectum, Open Approach		1	
48.59 Other abdominoperineal resection of the rectum	0DTP7ZZ Resection of Rectum, Via Natural or Artificial Opening; 0DTP8ZZ Resection of Rectum, Via Natural or Artificial Opening Endoscopic		2	
53.42 Laparoscopic repair of umbilical hernia with graft or prosthesis	0WUF4[7,J-K]Z Supplement Abdominal Wall with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 2 others	3		[Char 3] U Supplement

New ICD-9-CM	ICD-10-PCS	# New PCS Codes	# PCS Codes	New PCS Value
53.43 Other laparoscopic umbilical herniorrhaphy	0WQF4ZZ Repair Abdominal Wall, Percutaneous Endoscopic Approach		1	
53.62 Laparoscopic incisional hernia repair with graft or prosthesis	0WUF4[7,J-K]Z Supplement Abdominal Wall with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 2 others	3		[Char 3] U Supplement
53.63 Other laparoscopic repair of other hernia of anterior abdominal wall with graft or prosthesis	0WUF4[7,J-K]Z Supplement Abdominal Wall with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 2 others	3		[Char 3] U Supplement
53.71 Laparoscopic repair of diaphragmatic hernia, abdominal approach	0BU[R,S]4[7,J-K]Z ; 0BQ[R,S]4ZZ <u>Supplement Right Diaphragm</u> with <u>Autologous Tissue Substitute</u> , Percutaneous Endoscopic Approach + 7 others	6	2	[Char 3] U Supplement
53.72 Other and open repair of diaphragmatic hernia, abdominal approach	0BU[R,S]0[7,J-K]Z; 0BQ[R,S][0,3]ZZ <u>Supplement Right Diaphragm</u> with <u>Autologous Tissue Substitute</u> , Open Approach + 9 others	6	4	[Char 3] U Supplement
53.75 Repair of diaphragmatic hernia, abdominal approach, not otherwise specified	0BU[R,S][0,4][7,J-K]Z; 0BQ[R,S][0,3-4]ZZ <u>Supplement Right Diaphragm</u> with <u>Autologous Tissue Substitute</u> , Open Approach + 17 others	12	6	[Char 3] U Supplement
53.83 Laparoscopic repair of diaphragmatic hernia, with thoracic approach	0BQR4ZZ Repair Right Diaphragm, Percutaneous Endoscopic Approach; 0BQS4ZZ Repair Left Diaphragm, Percutaneous Endoscopic Approach		2	
53.84 Other and open repair of diaphragmatic hernia, with thoracic approach	0BQ[R,S][0,4]ZZ Repair <u>Right Diaphragm</u> , <u>Open</u> Approach + 3 others		4	
80.53 Repair of the anulus fibrosus with graft or prosthesis	0RU[3,9,B]0[7,J,K]Z; 0SU[2,4]0[7,J,K]Z <u>Supplement Cervical Vertebral Disc</u> with <u>Autologous Tissue Substitute</u> , Open Approach + 14 others	15		[Char 3] U Supplement
80.54 Other and unspecified repair of the anulus fibrosus	0RQ[3,9,B]0ZZ ; 0SQ[2,4]0ZZ <u>Repair Cervical Vertebral Disc</u> , Open Approach + 4 others		5	
85.70 Total reconstruction of breast, not otherwise specified	0HR[T-V]07Z Replacement of <u>Right Breast</u> with <u>Autologous Tissue Substitute</u> , Open Approach + 5 others		6	
85.71 Latissimus dorsi myocutaneous flap	0HR[T-V]075 Replacement of <u>Right Breast</u> using Latissimus Dorsi Myocutaneous Flap, Open Approach + 2 others	3		[Char 7] 5 Latissimus Dorsi Myocutaneous Flap
85.72 Transverse rectus abdominis myocutaneous (TRAM) flap, pedicled	0KX[K-L][0,4]Z6 Transfer <u>Right Abdomen Muscle</u> , Transverse Rectus Abdominis Myocutaneous Flap, <u>Open</u> Approach + 3 others	4		[Char 7] 6 Transverse Rectus Abdominis Myocutaneous Flap
85.73 Transverse rectus abdominis myocutaneous (TRAM) flap, free	0HR[T-V]076 Replacement of <u>Right Breast</u> using Transverse Rectus Abdominis Myocutaneous Flap, Open Approach + 2 others	3		[Char 7] 6 Transverse Rectus Abdominis Myocutaneous Flap
85.74 Deep inferior epigastric artery perforator (DIEP) flap, free	0HR[T-V]077 Replacement of <u>Right Breast</u> using Deep Inferior Epigastric Artery Perforator Flap, Open Approach + 2 others	3		[Char 7] 7 Deep Inferior Epigastric Artery Perforator Flap

New ICD-9-CM	ICD-10-PCS	# New PCS Codes	# PCS Codes	New PCS Value
85.75 Superficial inferior epigastric artery (SIEA) flap, free	0HR[T-V]078 Replacement of <u>Right Breast</u> using Superficial Inferior Epigastric Artery Flap, Open Approach + 2 others	3		[Char 7] 8 Superficial Inferior Epigastric Artery Flap
85.76 Gluteal artery perforator (GAP) flap, free	0HR[T-V]079 Replacement of <u>Right Breast</u> using Gluteal Artery Perforator Flap, Open Approach + 2 others	3		[Char 7] 9 Gluteal Artery Perforator Flap
85.79 Other total reconstruction of breast	0HR[T-V]07Z Replacement of <u>Right Breast</u> with <u>Autologous Tissue Substitute</u> , Open Approach + 5 others		6	