

Ambulance Fee Schedule - Medical Conditions List

Emergency Conditions - Non-Traumatic

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
535.50	458.9, 780.2, 787.01, 787.02, 787.03, 789.01, 789.02, 789.03, 789.04, 789.05, 789.06, 789.07, 789.09, 789.60 through 789.69, or 789.40 through 789.49 PLUS any other code from 780 through 799 except 793, 794, and 795.	Severe abdominal pain	With other signs or symptoms	ALS	Nausea, vomiting, fainting, pulsatile mass, distention, rigid, tenderness on exam, guarding.	A0427/A0433
789.00	726.2, 789.01, 789.02, 789.03, 789.04, 789.05, 789.06, 789.07, or 789.09	Abdominal pain	Without other signs or symptoms	BLS		A0429
427.9	426.0, 426.3, 426.4, 426.6, 426.11, 426.13, 426.50, 426.53, 427.0, 427.1, 427.2, 427.31, 427.32, 427.41, 427.42, 427.5, 427.60, 427.61, 427.69, 427.81, 427.89, 785.0, 785.50, 785.51, 785.52, or 785.59.	Abnormal cardiac rhythm/Cardiac dysrhythmia.	Potentially life-threatening	ALS	Bradycardia, junctional and ventricular blocks, non-sinus tachycardias, PVC's >6, bi- and trigeminy, ventricular tachycardia, ventricular fibrillation, atrial flutter, PEA, asystole, AICD/AED fired	A0427/A0433
780.8	782.5 or 782.6	Abnormal skin signs		ALS	Diaphoresis, cyanosis, delayed cap refill, poor turgor, mottled.	A0427/A0433
796.4	458.9, 780.6, 785.9, 796.2, or 796.3 PLUS any other code from 780 through 799	Abnormal vital signs (includes abnormal pulse oximetry)	With or without symptoms.	ALS		A0427/A0433
995.0	995.1, 995.2, 995.3, 995.4, 995.60, 995.61, 995.62, 995.63, 995.64, 995.65, 995.66, 995.67, 995.68, 995.69, or 995.7	Allergic reaction	Potentially life-threatening	ALS	Other emergency conditions, rapid progression of symptoms, prior history of anaphylaxis, wheezing, difficulty swallowing.	A0427/A0433
692.9	692.0, 692.1, 692.2, 692.3, 692.4, 692.5, 692.6, 692.70, 692.71, 692.72, 692.73, 692.74, 692.75, 692.76, 692.77, 692.79, 692.81, 692.82, 692.83, 692.89, 692.9, 693.0, 693.1, 693.8, 693.9, 695.9, 698.9, 708.9, 782.1.	Allergic reaction	Other	BLS	Hives, itching, rash, slow onset, local swelling, redness, erythema.	A0429
790.21	790.22, 250.02, or 250.03.	Blood glucose	Abnormal <80 or >250, with symptoms.	ALS	Altered mental status, vomiting, signs of dehydration.	A0427/A0433

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
799.1	786.02, 786.03, 786.04, or 786.09.	Respiratory arrest		ALS	Apnea, hypoventilation requiring ventilatory assistance and airway management.	A0427/A0433
786.05		Difficulty breathing		ALS		A0427/A0433
427.5		Cardiac arrest - resuscitation in progress		ALS		A0427/A0433
786.50	786.51, 786.52, or 786.59.	Chest pain (non-traumatic)		ALS	Dull, severe, crushing, substernal, epigastric, left sided chest pain associated with pain of the jaw, left arm, neck, back, and nausea, vomiting, palpitations, pallor, diaphoresis, decreased LOC.	A0427/A0433
784.99	933.0 or 933.1.	Chocking episode	Airway obstructed or partially obstructed	ALS		A0427/A0433
991.6		Cold exposure	Potentially life or limb threatening	ALS	Temperature < 95F, deep frost bite, other emergency conditions.	A0427/A0433
991.9	991.0, 991.1, 991.2, 991.3, or 991.4.	Cold exposure	With symptoms	BLS	Shivering, superficial frost bite, and other emergency conditions	A0429
780.97	780.02, 780.03, or 780.09.	Altered level of consciousness (nontraumatic)		ALS	Acute condition with Glasgow Coma Scale < 15.	A0427/A0433
780.39	345.00, 345.01, 345.2, 345.3, 345.10, 345.11, 345.40, 345.41, 345.50, 345.51, 345.60, 345.61, 345.70, 345.71, 345.80, 345.81, 345.90, 345.91, or 780.31.	Convulsions, seizures	Seizing, immediate post-seizure, postictal, or at risk of seizure and requires medical monitoring/observation.	ALS		A0427/A0433
379.90	368.11, 368.12, or 379.91	Eye symptoms, non-traumatic	Acute vision loss and/or severe pain	BLS		A0429
437.9	784.0 PLUS 781.0, 781.1, 781.2, 781.3, 781.4, or 781.8.	Non-traumatic headache	With neurologic distress conditions or sudden severe onset	ALS		A0427/A0433
785.1		Cardiac symptoms other than chest pain.	Palpitations, skipped beats	ALS		A0472/A0433

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
536.2	787.01, 787.02, 787.03, 780.79, 786.8, or 786.52.	Cardiac symptoms other than chest pain.	Atypical pain or other symptoms	ALS	Persistent nausea and vomiting, weakness, hiccups, pleuritic pain, feeling of impending doom, and other emergency conditions.	A0427/A0433
992.5	992.0, 992.1, 992.3, 992.4, or 992.5	Heat exposure	Potentially life-threatening	ALS	Hot and dry skin, Temp>105, neurologic distress, signs of heat stroke or heat exhaustion, orthostatic vitals, other emergency conditions.	A0427/A0433
992.2	992.6, 992.7, 992.8, or 992.9.	Heat exposure	With symptoms	BLS	Muscle cramps, profuse sweating, fatigue.	A0429
459.0	569.3, 578.0, 578.1, 578.9, 596.7, 596.8, 623.8, 626.9, 637.1, 634.1, 666.00, 666.02, 666.04, 666.10, 666.12, 666.14, 666.20, 666.22, 666.24, 674.30, 674.32, 674.34, 786.3, 784.7, or 998.11	Hemorrhage	Severe (quantity) and potentially life-threatening	ALS	Uncontrolled or significant signs of shock or other emergency conditions. Severe, active vaginal, rectal bleeding, hematemesis, hemoptysis, epistaxis, active post- surgical bleeding.	A0472/A0433
038.9	136.9, any other condition in the 001 through 139 code range which would require isolation.	Infectious diseases requiring isolation procedures / public health risk.		BLS		A0429
987.9	981, 982.0, 982.1, 982.2, 982.3, 982.4, 982.8, 983.0, 983.1, 983.2, 983.9, 984.0, 984.1, 984.8, 984.9, 985.0, 985.1, 985.2, 985.3, 985.4, 985.5, 985.6, 985.8, 985.9, 986, 987.0, 987.1, 987.2, 987.3, 987.4, 987.5, 987.6, 987.7, 987.8, 989.1, 989.2, 989.3, 989.4, 989.6, 989.7, 989.9, or 990.	Hazmat exposure		ALS	Toxic fume or liquid exposure via inhalation, absorption, oral, radiation, smoke inhalation.	A0472/A0433
996.00	996.01, 996.02, 996.04, 996.09, 996.1, or 996.2.	Medical device failure	Life or limb threatening malfunction, failure, or complication.	ALS	Malfunction of ventilator, internal pacemaker, internal defibrillator, implanted drug delivery service.	A0427/A0433
996.30	996.31, 996.40, 996.41, 996.42, 996.43, 996.44, 996.45, 996.46, 996.47, 996.49, or 996.59.	Medical device failure	Health maintenance device failures that cannot be resolved on location.	BLS	Oxygen system supply malfunction, orthopedic device failure.	A0429

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
436	291.3, 293.82, 298.9, 344.9, 368.16, 369.9, 780.09, 780.4, 781.0, 781.2, 781.94, 781.99, 782.0, 784.3, 784.5, or 787.2.	Neurologic distress	Facial drooping; loss of vision; aphasia; difficulty swallowing; numbness, tingling extremity; stupor, delirium, confusion, hallucinations; paralysis, paresis (focal weakness); abnormal movements; vertigo; unsteady gait/ balance; slurred speech, unable to speak	ALS		A0427/A0433
780.96		Pain, severe not otherwise specified in this list.	Acute onset, unable to ambulate or sit due to intensity of pain.	ALS	Pain is the reason for the transport. Use severity scale (7-10 for severe pain) or patient receiving pharmacologic intervention.	A0427/A0433
724.5	724.2 or 785.9	Back pain - non-traumatic (T and/or LS).	Suspect cardiac or vascular etiology	ALS	Other emergency conditions, absence of or decreased leg pulses, pulsatile abdominal mass, severe tearing abdominal pain.	A0427/A0433
724.9	724.2, 724.5, 847.1, or 847.2.	Back pain - non-traumatic (T and/or LS).	Sudden onset of new neurologic symptoms.	ALS	Neurologic distress list.	A0427/A0433
977.9	Any code from 960 through 979.	Poisons, ingested, injected, inhaled, absorbed.	Adverse drug reaction, poison exposure by inhalation, injection, or absorption.	ALS		A0427/A0433
305.0	303.00, 303.01, 303.02, 303.03, or any code from 960 through 979.	Alcohol intoxication or drug overdose (suspected).	Unable to care for self and unable to ambulate. No airway compromise.	BLS		A0429

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
977.3		Severe alcohol intoxication.	Airway may or may not be at risk. Pharmacological intervention or cardiac monitoring may be needed. Decreased level of consciousness resulting or potentially resulting in airway compromise.	ALS		A0427/A0433
998.9	674.10, 674.12, 674.14, 674.20, 674.22, 674.24, 997.69, 998.31, 998.32, or 998.83.	Post-operative procedure complications.	Major wound dehiscence, evisceration, or requires special handling for transport.	BLS	Non-life threatening	A0429
650	Any code from 630 through 679.	Pregnancy complication/childbirth/labor		ALS		A0427/A0433
292.9	291.0, 291.3, 291.81, 292.0, 292.81, 292.82, 292.83, 292.84, or 292.89.	Psychiatric/Behavioral	Abnormal mental status; drug withdrawal.	ALS	Disoriented, DTs, withdrawal symptoms.	A0427/A0433
298.9	300.9	Psychiatric/Behavioral	Threat to self or others, acute episode or exacerbation of paranoia, or disruptive behavior.	BLS	Suicidal, homicidal, or violent.	A0429
036.9	780.6 PLUS either 784.0 or 723.5.	Sick person - fever	Fever with associated symptoms (headache, stiff neck, etc.). Neurological changes.	BLS	Suspected spinal meningitis.	A0429

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
787.01	787.02, 787.03, or 787.91.	Severe dehydration	Nausea and vomiting, diarrhea, severe and incapacitating resulting in severe side effects of dehydration.	ALS		A0427/A0433
780.02	780.2 or 780.4	Unconscious, fainting, syncope, near syncope, weakness, or dizziness.	Transient unconscious episode or found unconscious. Acute episode or exacerbation.	ALS		A0427/A0433

Emergency Conditions - Trauma

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
959.8	800.00 through 804.99, 807.4, 807.6, 808.8, 808.9, 812.00 through 812.59, 813.00 through 813.93, 813.93, 820.00 through 821.39, 823.00 through 823.92, 851.00 through 866.13, 870.0 through 879.9, 880.00 through 887.7, or 890.0 through 897.7.	Major trauma	As defined by ACS Field Triage Decision Scheme. Trauma with one of the following: Glasgow <14; systolic BP<90; RR<10 or >29; all penetrating injuries to head, neck, torso, extremities proximal to elbow or knee; flail chest; combination of trauma and burns; pelvic fracture; 2 or more long bone fractures; open or depressed skull fracture; paralysis; severe mechanism of injury including: ejection, death of another passenger in same patient compartment, falls >20", 20" deformity in vehicle or 12" deformity of patient compartment, auto pedestrian/bike, pedestrian thrown/run over, motorcycle accident at speeds >20 mph and rider separated from vehicle.	ALS	See "Condition (Specific)" column	A0427/A0433

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
518.5		Other trauma	Need to monitor or maintain airway	ALS	Decreased LOC, bleeding into airway, trauma to head, face or neck.	A0427/A0433
958.2	870.0 through 879.9, 880.00 through 887.7, 890.0 through 897.7, or 900.00 through 904.9.	Other trauma	Major bleeding	ALS	Uncontrolled or significant bleeding.	A0427/A0433
829.0	805.00, 810.00 through 819.1, or 820.00 through 829.1.	Other trauma	Suspected fracture/dislocation requiring splinting/immobilization for transport.	BLS	Spinal, long bones, and joints including shoulder elbow, wrist, hip, knee and ankle, deformity of bone or joint.	A0429
880.00	880.00 through 887.7 or 890.0 through 897.7	Other trauma	Penetrating extremity injuries	BLS	Isolated bleeding stopped and good CSM.	A0429
886.0 or 895.0	886.1 or 895.1	Other trauma	Amputation - digits	BLS		A0429
887.4 or 897.4	887.0, 887.1, 887.2, 887.3, 887.6, 887.7, 897.0, 897.1, 897.2, 897.3, 897.5, 897.6, or 897.7.	Other trauma	Amputation - all other	ALS		A0427/A0433
869.0 or 869.1	511.8, 512.8, 860.2, 860.3, 860.4, 860.5, 873.8, 873.9, or 959.01.	Other trauma	Suspected internal, head, chest, or abdominal injuries.	ALS	Signs of closed head injury, open head injury, pneumothorax, hemothorax, abdominal bruising, positive abdominal signs on exam, internal bleeding criteria, evisceration.	A0427/A0433

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
949.3	941.30 through 941.39, 942.30 through 942.39, 943.30 through 943.39, 944.30 through 944.38, 945.30 through 945.39, or 949.3.	Burns	Major - per American Burn Association (ABA)	ALS	Partial thickness burns > 10% total body surface area (TBSA); involvement of face, hands, feet, genitalia, perineum, or major joints; third degree burns; electrical; chemical; inhalation; burns with preexisting medical disorders; burns and trauma	A0472/A0433
949.2	941.20 through 941.29, 942.20 through 942.29, 943.20 through 943.29, 944.20 through 944.28, 945.20 through 945.29, or 949.2.	Burns	Minor - per ABA	BLS	Other burns than listed above.	A0429
989.5		Animal bites, stings, envenomation.	Potentially life or limb-threatening.	ALS	Symptoms of specific envenomation, significant face, neck, trunk, and extremity involvement; other emergency conditions.	A0427/A0433
879.8	Any code from 870.0 through 897.7.	Animal bites/sting/envenomation.	Other	BLS	Local pain and swelling or special handling considerations (not related to obesity) and patient monitoring required.	A0429
994.0		Lightning		ALS		A0427/A0433
994.8		Electrocution		ALS		A0427/A0433

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
994.1		Near drowning	Airway compromised during near drowning event	ALS		A0427/A0433
921.9	870.0 through 870.9, 871.0, 871.1, 871.2, 871.3, 871.4, 871.5, 871.6, 871.7, 871.9, or 921.0 through 921.9.	Eye injuries	Acute vision loss or blurring, severe pain or chemical exposure, penetrating, severe lid lacerations.	BLS		A0429
995.83	995.53 or V71.5 PLUS any code from 925.1 through 929.9, 930.0 through 939.9, 958.0 through 958.8, or 959.01 through 959.9.	Sexual assault	With major injuries	ALS	Reference codes 959.8, 958.2, 869.0/869.1	A0427/A0433
995.80	995.53 or V71.5 PLUS any code from 910.0 through 919.9, 920 through 924.9, or 959.01 through 959.9.	Sexual assault	With minor or no injuries	BLS		

Non-Emergency

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
428.9		Cardiac/hemodynamic monitoring required en route.		ALS	Expectation monitoring is needed before and after transport.	A0426
518.81 or 518.89	V46.11 or V46.12.	Advanced airway management		ALS	Ventilator dependent, apnea monitor, possible intubation needed, deep suctioning.	A0426, A0434
293.0		Chemical restraint.		ALS		A0426
496	491.20, 491.21, 492.0 through 492.8, 493.20, 493.21, 493.22, 494.0, or 494.1.	Suctioning required en route, need for titrated O2 therapy or IV fluid management.		BLS	Per transfer instructions.	A0428
786.09		Airway control/positioning required en route.		BLS	Per transfer instructions.	A0428
492.8	491.20, 491.21, 492.0 through 492.8, 493.20, 493.21, 493.22, 494.0, or 494.1.	Third party assistance/attendant required to apply, administer, or regulate or adjust oxygen en route.		BLS	Does not apply to patient capable of self-administration of portable or home O2. Patient must require oxygen therapy and be so frail as to require assistance.	A0428
298.9	Add 295.0 through 295.9 with 5 th digits of 0, 1, 3, or 4, 296.00 or 299.90.	Patient safety: Danger to self or others - in restraints.		BLS	Refer to definition in 42 CFR Section 482.13(e).	A0428
293.1		Patient safety: Danger to self or others - monitoring.		BLS	Behavioral or cognitive risk such that patient requires monitoring for safety.	A0428

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
298.8	Add 295.0 through 295.9 with 5 th digits of 0, 1, 3, or 4, 296.00 or 299.90	Patient safety: Danger to self or others - seclusion (flight risk).		BLS	Behavioral or cognitive risk such that patient requires attendant to assure patient does not try to exit the ambulance prematurely. Refer to 42 CFR Section 482.13(f) for definition.	A0428
781.3	Add 295.0 through 295.9 with 5 th digits of 0, 1, 3, or 4, 296.00 or 299.90.	Patient safety: Risk of falling off wheelchair or stretcher while in motion (not related to obesity).		BLS	Patient's physical condition is such that patient risks injury during vehicle movement despite restraints. Indirect indicators include MDS criteria.	A0428
041.9		Special handling en route - isolation.		BLS	Includes patients with communicable diseases or hazardous material exposure who must be isolated from public or whose medical condition must be protected from public exposure; surgical drainage complications.	A0428
907.2		Special handling en route to reduce pain - orthopedic device.		BLS	Backboard, halotraction, use of pins and traction etc. Pain may be present.	A0428

ICD 9 Primary Code	ICD9 Alternative Specific Code	Condition (General)	Condition (Specific)	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
719.45 or 719.49	718.40, 718.45, 718.49, or 907.2.	Special handling en route - positioning requires specialized handling.		BLS	Requires special handling to avoid further injury (such as with > grade 2 decubiti on buttocks). Generally does not apply to shorter transfers of < 1 hour. Positioning in wheelchair or standard car seat inappropriate due to contractures or recent extremity fractures - post-op hip as an example.	A0428

Transportation Indicators

Transportation Indicators Air and Ground	Transportation Category	Transportation Indicator Description		Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
C1	Inter-facility Transport	EMTALA-certified inter-facility transfer to a higher level of care.	Beneficiary requires higher level of care.	BLS, ALS, SCT, FW, RW	Excludes patient-requested EMTALA transfer.	A0428, A0429, A0426, A0427, A0433, A0434
C2	Inter-facility transport	Service not available at originating facility, and must meet one or more emergency or non-emergency conditions.		BLS, ALS, SCT, FW, RW		A0428, A0429, A0426, A0427, A0433, A0434
C3	Emergency Trauma Dispatch Condition Code	Major incident or mechanism of injury	Major Incident-This transportation indicator is to be used ONLY as a secondary code when the on-scene encounter is a BLS-level patient.	ALS	Trapped in machinery, close proximity to explosion, building fire with persons reported inside, major incident involving aircraft, bus, subway, metro, train and watercraft. Victim entrapped in vehicle.	A0427/A0433
C4	Medically necessary transport but not to the nearest facility.	BLS or ALS response	Indicates to A/B MAC (A or B) that an ambulance provided a medically necessary transport, but that the number of miles on the Medicare claim form may be excessive.	BLS/ALS	This should occur if the facility is on divert status or the particular service is not available at the time of transport only. In these instances the ambulance units should clearly document why the beneficiary was not transported to the nearest facility.	Based on transport level.

Transportation Indicators Air and Ground	Transportation Category	Transportation Indicator Description		Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
C5	BLS transport of ALS-level patient	ALS-level condition treated and transport by a BLS-level ambulance.	This transportation indicator is used for ALL situations where a BLS-level ambulance treats and transports a patient that presents an ALS-level condition. No ALS-level assessment or intervention occurs at all during the patient encounter.	BLS		A0429
C6	ALS-level response to BLS-level patient	ALS response required based upon appropriate dispatch protocols - BLS-level patient transport	Indicates to A/B MAC (A or B) that an ALS-level ambulance responded appropriately based upon the information received at the time the call was received in dispatch and after a clinically appropriate ALS-assessment was performed on scene, it was determined that the condition of the patient was at a BLS level. These claims, properly documented, should be reimbursed at an ALS-1 level based upon coverage guidelines under the Medicare Ambulance Fee Schedule.	ALS		A0427

Transportation Indicators Air and Ground	Transportation Category	Transportation Indicator Description		Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
C7		IV meds required en route.	This transportation indicator is used for patients that require an ALS level transport in a non-emergent situation primarily because the patient requires monitoring of ongoing medications administered intravenously. Does not apply to self-administered medications. Does not include administration of crystalloid intravenous fluids (i.e., Normal Saline, Lactate Ringers, 5% Dextrose in Water, etc.). The patient's condition should also be reported on the claim with a code selected from the list	ALS	Does not apply to self-administered IV medications.	A0426

Air Ambulance Transportation Indicators

Air Ambulance Transportation Indicators	Transportation Indicator Description	Service Level	Comments and Examples (not all-inclusive)	HCPCS Crosswalk
D1	Long Distance-patient's condition requires rapid transportation over a long distance	FW, RW	If the patient's condition warrants only.	A0430, A0431
D2	Under rare and exceptional circumstances, traffic patterns preclude ground transport at the time the response is required.	FW, RW		A0430, A0431
D3		FW, RW		A0430, A0431
D4		FW, RW		A0430, A0431

Note: HCPCS Crosswalk to ALS1E (A0427) and ALS2 (A0433) would ultimately be determined by the number and type of ALS level services provided during transport. All medical condition codes can be cross walked to fixed wing and rotor wing HCPCS provided the air ambulance service has documented the medical necessity for air ambulance service versus ground or water ambulance. As a result, codes A0430 (Fixed Wing) and A0431 (Rotor Wing) can be included in Column 7 for each condition listed.